

Final Size : 290 (L) X 185 (H) mm



EMAMECTIN BENZOATE 5% SG (Insecticide)

Emamectin Benzoate 5% SG is a water soluble granular formulation containing 5% active ingredient and is recommended for the control of blowfly on cotton, fruit and vegetable crops... [Detailed text about usage and efficacy]

Table with 5 columns: Crop, Area (ha), Recommended Rate (g/ha), and other parameters.

नॅटजोएट इमामेक्टिन बॅन्झोएट ५% एस. जी.

Marketed by : Natco Pharma Limited, Natco House, Road No. 2, Banjara Hills, Hyderabad, Telangana- 500034

Mfg. By: Crystal Crop Protection Ltd. Factory Unit (A) Lane No.3 (C), Phase-II, SIDCO Industrial Complex, Bari-Brahmana, Jammu- 181133 (J&K)...

इमामेक्टिन बॅन्झोएट ५% एस. जी.

इमामेक्टिन बॅन्झोएट ५% एस. जी. एक जलघुलनशील कणिकार प्रारूप में 5% सक्रिय घटक युक्त प्रारूप है... [Detailed text about usage and efficacy]

Table with 5 columns: Crop, Area (ha), Recommended Rate (g/ha), and other parameters.

उपरोक्त सूची में उल्लेखित सभी फसलों पर इस कीटनाशक का प्रयोग करने के लिए निर्देशों का पालन करना चाहिए... [Detailed text about application and safety]

Emamectin Benzoate 5% SG

Emamectin Benzoate 5% SG is a water soluble granular formulation containing 5% active ingredient... [Detailed text about usage and efficacy]

Table with 5 columns: Crop, Area (ha), Recommended Rate (g/ha), and other parameters.

This insecticide is used for the control of blowfly on cotton, fruit and vegetable crops... [Detailed text about application and safety]

इमामेक्टिन बॅन्झोएट ५% एस. जी.

इमामेक्टिन बॅन्झोएट ५% एस. जी. एक जलघुलनशील कणिकार प्रारूप में 5% सक्रिय घटक युक्त प्रारूप है... [Detailed text about usage and efficacy]

Table with 5 columns: Crop, Area (ha), Recommended Rate (g/ha), and other parameters.

इमामेक्टिन बॅन्झोएट ५% एस. जी. का प्रयोग करने के लिए निर्देशों का पालन करना चाहिए... [Detailed text about application and safety]

Emamectin Benzoate 5% SG

Emamectin Benzoate 5% SG is a water soluble granular formulation containing 5% active ingredient... [Detailed text about usage and efficacy]

Table with 5 columns: Crop, Area (ha), Recommended Rate (g/ha), and other parameters.

This insecticide is used for the control of blowfly on cotton, fruit and vegetable crops... [Detailed text about application and safety]

Emamectin Benzoate 5% SG

Emamectin Benzoate 5% SG is a water soluble granular formulation containing 5% active ingredient... [Detailed text about usage and efficacy]

Table with 5 columns: Crop, Area (ha), Recommended Rate (g/ha), and other parameters.

This insecticide is used for the control of blowfly on cotton, fruit and vegetable crops... [Detailed text about application and safety]

EMPTYPACKAGING SHOULD BE DESTROYED AFTER USE


READ THE ENTIRE LABEL AND LEAFLET BEFORE USE. FOLLOW INSTRUCTIONS CAREFULLY. Empty packaging should be destroyed after use.



For Consumer Complaints / Feedback-Customer Care Executive. Contact details and email address.

EMPTYPACKAGING SHOULD BE DESTROYED AFTER USE


READ THE ENTIRE LABEL AND LEAFLET BEFORE USE. FOLLOW INSTRUCTIONS CAREFULLY. Empty packaging should be destroyed after use.



For Consumer Complaints / Feedback-Customer Care Executive. Contact details and email address.

EMPTYPACKAGING SHOULD BE DESTROYED AFTER USE


READ THE ENTIRE LABEL AND LEAFLET BEFORE USE. FOLLOW INSTRUCTIONS CAREFULLY. Empty packaging should be destroyed after use.



For Consumer Complaints / Feedback-Customer Care Executive. Contact details and email address.

EMPTYPACKAGING SHOULD BE DESTROYED AFTER USE


READ THE ENTIRE LABEL AND LEAFLET BEFORE USE. FOLLOW INSTRUCTIONS CAREFULLY. Empty packaging should be destroyed after use.



For Consumer Complaints / Feedback-Customer Care Executive. Contact details and email address.

EMPTYPACKAGING SHOULD BE DESTROYED AFTER USE


READ THE ENTIRE LABEL AND LEAFLET BEFORE USE. FOLLOW INSTRUCTIONS CAREFULLY. Empty packaging should be destroyed after use.



For Consumer Complaints / Feedback-Customer Care Executive. Contact details and email address.

EMPTYPACKAGING SHOULD BE DESTROYED AFTER USE

READ THE ENTIRE LABEL AND LEAFLET BEFORE USE. FOLLOW INSTRUCTIONS CAREFULLY. Empty packaging should be destroyed after use.



For Consumer Complaints / Feedback-Customer Care Executive. Contact details and email address.



EMAMECTIN BENZOATE 5% SG (Insecticide)

Emamectin Benzoate 5% SG is a water soluble granular formulation containing 5% active ingredient and is recommended for the control of blowfly on cotton, fruit and vegetable crops...

Table with 5 columns: Crop, Area (ha), Recommended Rate (g/ha), and other parameters.

नॅटजोएट इमामेक्टिन बॅन्झोएट ५% एस. जी.

Marketed by : Natco Pharma Limited, Natco House, Road No. 2, Banjara Hills, Hyderabad, Telangana- 500034

Mfg. By: Crystal Crop Protection Ltd. Factory Unit (A) Lane No.3 (C), Phase-II, SIDCO Industrial Complex, Bari-Brahmana, Jammu- 181133 (J&K)...

For Consumer Complaints / Feedback-Customer Care Executive. Contact details and email address.

इमामेक्टिन बॅन्झोएट ५% एस. जी.

इमामेक्टिन बॅन्झोएट ५% एस. जी. एक जलघुलनशील कणिकार प्रारूप में 5% सक्रिय घटक युक्त प्रारूप है... [Detailed text about usage and efficacy]

Table with 5 columns: Crop, Area (ha), Recommended Rate (g/ha), and other parameters.

इमामेक्टिन बॅन्झोएट ५% एस. जी.

इमामेक्टिन बॅन्झोएट ५% एस. जी. का प्रयोग करने के लिए निर्देशों का पालन करना चाहिए... [Detailed text about application and safety]

इमामेक्टिन बॅन्झोएट ५% एस. जी. का प्रयोग करने के लिए निर्देशों का पालन करना चाहिए... [Detailed text about application and safety]

Emamectin Benzoate 5% SG

Emamectin Benzoate 5% SG is a water soluble granular formulation containing 5% active ingredient... [Detailed text about usage and efficacy]

Table with 5 columns: Crop, Area (ha), Recommended Rate (g/ha), and other parameters.

Emamectin Benzoate 5% SG

This insecticide is used for the control of blowfly on cotton, fruit and vegetable crops... [Detailed text about application and safety]

This insecticide is used for the control of blowfly on cotton, fruit and vegetable crops... [Detailed text about application and safety]

इमामेक्टिन बॅन्झोएट ५% एस. जी.

इमामेक्टिन बॅन्झोएट ५% एस. जी. एक जलघुलनशील कणिकार प्रारूप में 5% सक्रिय घटक युक्त प्रारूप है... [Detailed text about usage and efficacy]

Table with 5 columns: Crop, Area (ha), Recommended Rate (g/ha), and other parameters.

इमामेक्टिन बॅन्झोएट ५% एस. जी.

इमामेक्टिन बॅन्झोएट ५% एस. जी. का प्रयोग करने के लिए निर्देशों का पालन करना चाहिए... [Detailed text about application and safety]

इमामेक्टिन बॅन्झोएट ५% एस. जी. का प्रयोग करने के लिए निर्देशों का पालन करना चाहिए... [Detailed text about application and safety]

Emamectin Benzoate 5% SG

Emamectin Benzoate 5% SG is a water soluble granular formulation containing 5% active ingredient... [Detailed text about usage and efficacy]

Table with 5 columns: Crop, Area (ha), Recommended Rate (g/ha), and other parameters.

Emamectin Benzoate 5% SG

This insecticide is used for the control of blowfly on cotton, fruit and vegetable crops... [Detailed text about application and safety]

This insecticide is used for the control of blowfly on cotton, fruit and vegetable crops... [Detailed text about application and safety]

Emamectin Benzoate 5% SG

Emamectin Benzoate 5% SG is a water soluble granular formulation containing 5% active ingredient... [Detailed text about usage and efficacy]

Table with 5 columns: Crop, Area (ha), Recommended Rate (g/ha), and other parameters.

Emamectin Benzoate 5% SG

This insecticide is used for the control of blowfly on cotton, fruit and vegetable crops... [Detailed text about application and safety]

This insecticide is used for the control of blowfly on cotton, fruit and vegetable crops... [Detailed text about application and safety]